

BRAID GROUP OF A GENETIC CODE

VALERIJ G. BARDAKOV

ABSTRACT. For every group genetic code with finite number of generating and at most with one defining relation we introduce the braid group of this genetic code. This construction includes the braid group of Euclidean plane, the braid groups of closed orientable surfaces, B type groups of Artin-Brieskorn, and allow us to study all these groups with common point of view. Then we clarify a structure of the braid group of genetic code, construct a normal form of words, investigate a torsion, calculate the width of verbal subgroups and establish some other properties. Also we prove that Scott's system of defining relations for the braid groups of closed surfaces is contradictory.

In last years appeared many articles dealing with the braid groups and their different generalizations [1]-[9]. This fact is explained by connection of theory of braid groups with knot theory. Knot theory draws attention of many researchers with connection to appearance of new methods and directions, which arise in this theory. It is enough to remember about results of Jones [10], who constructed new knot polynomial invariant, about results of V. A. Vasilieva [3], who constructed finite type invariants, about results of Bigelow and Krammer [11, 12], who proved that the braid groups are linear, and also about result of Dehornoy [13] which said that the braid groups are right orderable. Apart from that knot theory is fast connects with other areas of mathematics and physics [14, 15].

One of the first generalizations of the braid groups are braid groups of closed orientable surfaces, which were introduced by Zariski [16]. Then in the Scott's article [17] were considered the braid groups of closed orientable surfaces and were found generators and defining relations of these groups and also of pure braid groups of these surfaces (a set of generators and defining relations is called the *genetic code* of this group). Unfortunately the set of defining relations which was written in the Scott's article is contradictory (see § 2 of the present paper). Braid groups of sphere and projecting plane differ from the braid groups of other surfaces, and as a rule are studied separately (see [18, 19, 20]).

González-Meneses [21] found a new presentation of the braid groups of closed surfaces in the form of a set of generators and defining relations. Then Bellingeri [22] simplified this presentation and found generators and defining relations of the braid groups of surfaces with boundary. In preprint [23] was found a positive presentation of the braid groups of orientable surfaces. Apart from that, in [21, 22] was solved the word problem for corresponding braid groups, and in preprint [23] was solved the conjugacy problem for the braid group $B_2(T^2)$ on two strands of torus T^2 . In general case the conjugacy problem for the braid groups of surfaces is still open.

Date: February 2, 2008.

1991 Mathematics Subject Classification. Primary 20F36; Secondary 20F05, 20F10.

Key words and phrases. Surface braid groups, group of a genetic code, width of verbal subgroups, free groups.

In the present paper we recall a definition of the braid groups of manifold (see § 1), show that Scott's presentation is contradictory (see § 2). As Bellingeri stated to the author that last fact was also observed by V. S. Kulikov and Shimada. Then in § 3 for each genetic code with finite number of generators and at most one relation we introduce the braid group of this genetic code. This construction includes the braid group of plane, the braid groups of closed orientable surfaces, the B -type of braid group of Artin-Briskorn, and it allows us to study all these groups with unified position. In § 4 we establish a structure of the braid group of genetic code, construct a normal form of words, investigate torsion, calculate width of verbal subgroups and establish some others properties.

The results of the present paper were announced in 2000 (see [24]), but at the series of reasons wasn't published in proper time.

Acknowledgments. I am very grateful to the participants of the seminar “Evariste Galois” who listened proofs and forwarded series of useful remarks and suggestions. Separately I thank M. V. Neshchadim for fruitful discussions and useful conversations. Also I thank V. V. Vershinin, who introduced Zariski's article to me. Special thanks goes to my brother Victor Bardakov who helped me to translate this article into English.

§ 1. Braid groups of manifolds

We recall some known facts about the braid groups of manifolds (see [25, 2]).

Let M be a connected manifold of dimension ≥ 2 . *Configuration space* $F_n M$, $n \in \mathbb{N}$, of manifold M is called the manifold

$$F_n M = \{(z_1, z_2, \dots, z_n) \in \prod_{i=1}^n M \mid z_i \neq z_j \text{ for } i \neq j\}$$

of ordering collections of n distinct points from M . Its fundamental group $\pi_1(F_n M)$ is called the n -strand pure braid group of manifold M and is denoted by $P_n(M)$. On the set $F_n M$ is defined an equivalence relations, $z \sim z'$ if the vector $z = (z_1, z_2, \dots, z_n) \in F_n M$ different from the vector $z' = (z'_1, z'_2, \dots, z'_n) \in F_n M$ by permutation components. The fundamental group $\pi_1(F_n M / \sim)$ of the factor manifold $F_n M / \sim$ is called the n -strand braid group of manifold M and denoted by $B_n(M)$. The natural projection $p : F_n M \longrightarrow F_n M / \sim$ is a regular covering projection. A group of covering transformations is the symmetric group S_n . Hence, there is a canonical isomorphism

$$\pi_1(F_n M / \sim) / \pi_1(F_n M) \simeq B_n(M) / P_n(M) \simeq S_n.$$

If M is a closed, smooth manifold then the map $F_n M \longrightarrow \prod_{i=1}^n M$ induces the epimorphism $P_n(M) \longrightarrow \pi_1(M) \times \dots \times \pi_1(M)$ of the pure braid group $P_n(M)$ on direct product n copies of fundamental group $\pi_1(M)$. And what is more, if the dimension of M is greater than two then this epimorphism is injective. Hence, the braid groups of manifolds of dimension 2 represent the largest interest. Further we will assume that M is a two dimension manifold. Braid group $B_n = B_n(E^2)$ of Euclidean plane E^2 is defined by generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ and defining relations

$$\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \text{ if } i = 1, 2, \dots, n-2,$$

$$\sigma_i \sigma_j = \sigma_j \sigma_i, \text{ if } |i - j| \geq 2.$$

The group $P_n = P_n(E^2)$ is generated by elements a_{ij} , $1 \leq i < j \leq n$, which express through the generators of B_n by the following manner:

$$a_{i,i+1} = \sigma_i^2,$$

$$a_{ij} = \sigma_{j-1}\sigma_{j-2}\dots\sigma_{i+1}\sigma_i^2\sigma_{i+1}^{-1}\dots\sigma_{j-2}^{-1}\sigma_{j-1}^{-1}, \quad i+1 < j \leq n.$$

For the pure braid group P_n there is an epimorphism $\eta : P_n \longrightarrow P_{n-1}$, its kernel $U_n = \ker(\eta)$ is a free group of rank $n-1$. The pure braid group P_n is a semi direct product of normal subgroup U_n , which is free generated by $a_{1,n}, a_{2,n}, \dots, a_{n-1,n}$ and the group P_{n-1} . Similar, P_{n-1} is a semi direct product of the free group U_{n-1} with free generators $a_{1,n-1}, a_{2,n-1}, \dots, a_{n-2,n-1}$ and the subgroup P_{n-2} , and so on. Hence, the group P_n has the following decomposition

$$P_n = U_n \rtimes (U_{n-1} \rtimes (\dots \rtimes (U_3 \rtimes U_2)) \dots), \quad U_i \simeq F_{i-1}, \quad i = 2, 3, \dots, n,$$

The pure braid group P_n is defined by the relations (for $\varepsilon = \pm$)

- 1) $a_{ik}^{-\varepsilon} a_{kj} a_{ik}^{\varepsilon} = (a_{ij} a_{kj})^{\varepsilon} a_{kj} (a_{ij} a_{kj})^{-\varepsilon},$
- 2) $a_{km}^{-\varepsilon} a_{kj} a_{km}^{\varepsilon} = (a_{kj} a_{mj})^{\varepsilon} a_{kj} (a_{kj} a_{mj})^{-\varepsilon}, \quad m < j,$
- 3) $a_{im}^{-\varepsilon} a_{kj} a_{im}^{\varepsilon} = [a_{ij}^{-\varepsilon}, a_{mj}^{-\varepsilon}]^{\varepsilon} a_{kj} [a_{ij}^{-\varepsilon}, a_{mj}^{-\varepsilon}]^{-\varepsilon}, \quad i < k < m,$
- 4) $a_{im}^{-\varepsilon} a_{kj} a_{im}^{\varepsilon} = a_{kj}, \quad k < i; \quad m < j \text{ or } m < k,$

where $[a, b] = a^{-1}b^{-1}ab$.

In this case generators of the braid group act on generators of the pure braid group by the formulas

- 1) $\sigma_k^{-\varepsilon} a_{ij} \sigma_k^{\varepsilon} = a_{ij}, \quad k \neq i-1, i, j-1, j,$
- 2) $\sigma_i^{-\varepsilon} a_{i,i+1} \sigma_i^{\varepsilon} = a_{i,i+1},$
- 3) $\sigma_{i-1}^{-1} a_{ij} \sigma_{i-1} = a_{i-1,j},$
- 4) $\sigma_{i-1} a_{ij} \sigma_{i-1}^{-1} = a_{i-1,i} a_{i-1,j} a_{i-1,i}^{-1},$
- 5) $\sigma_i^{-1} a_{ij} \sigma_i = a_{i+1,j} [a_{i,i+1}^{-1}, a_{ij}^{-1}], \quad j \neq i+1,$
- 6) $\sigma_i a_{ij} \sigma_i^{-1} = a_{i+1,j}, \quad j \neq i+1,$
- 7) $\sigma_{j-1}^{-1} a_{ij} \sigma_{j-1} = a_{i,j-1},$
- 8) $\sigma_{j-1} a_{ij} \sigma_{j-1}^{-1} = a_{ij}^{-1} a_{i,j-1} a_{ij},$
- 9) $\sigma_j^{-1} a_{ij} \sigma_j = a_{ij} a_{i,j+1} a_{ij}^{-1},$
- 10) $\sigma_j a_{ij} \sigma_j^{-1} = a_{i,j+1}, \quad 1 \leq i < j \leq n-1,$

(see [26]).

If $n \geq 3$ then the center of the group B_n is an infinite cyclic group, generated by an element

$$(\sigma_1 \sigma_2 \dots \sigma_{n-1})^n = a_{12}(a_{13}a_{23}) \dots (a_{1n}a_{2n} \dots a_{n-1,n}).$$

If M is a compact two dimension manifold which differs from the sphere S^2 and projective plane P^2 then the braid group $B_n(E^2)$ of Euclidean plane E^2 includes into the braid group $B_n(M)$ of manifold M for all $n \geq 1$.

Fadell and Noivirt proved that if M is a compact two dimension manifold which differs from S^2 and P^2 then there are not elements of finite order in the braid group $B_n(M)$.

The braid group $B_n(S^2)$ of sphere S^2 is generated by elements $\delta_1, \delta_2, \dots, \delta_{n-1}$ and defined by relations

$$\delta_i \delta_{i+1} \delta_i = \delta_{i+1} \delta_i \delta_{i+1} \text{ if } i = 1, 2, \dots, n-2,$$

$$\delta_i \delta_j = \delta_j \delta_i \text{ if } |i - j| \geq 2,$$

$$\delta_1 \delta_2 \dots \delta_{n-2} \delta_{n-1}^2 \delta_{n-2} \dots \delta_2 \delta_1 = 1.$$

In this case $P_2(S^2) = 1$, $P_3(S^2) = B_2(S^2) \simeq \mathbb{Z}_2$, $B_3(S^2)$ is a meta-cyclic group of order 12. If $n \geq 3$ then the center of $B_n(S^2)$ is a subgroup of order 2, generated by an element $(\delta_1 \delta_2 \dots \delta_{n-1})^n$. The group $B_{n-1}(S^2)$ is not included in the group $B_n(S^2)$ if $n \geq 3$.

§ 2. On Scott's presentation

If T_g is a compact orientable surface of genus $g \geq 1$ then generators and relations of the braid group $B_n(T_g)$, $n \geq 1$, was found in the work [17]. We will show that this system of defining relations is contradictorily.

Let $T = T_1$ be a compact orientable surface of genus 1 (torus). Let (US) consider the braid group $B_2(T)$ on two strands and its subgroup of the pure braid $P_2(T)$. As follows from Theorem 1.3 of Scott's paper, the group $P_2(T)$ is generated by elements

$$a_{12}, \rho_{11}, \rho_{12}, \rho_{21}, \rho_{22}$$

and defined by relations

$$[\rho_{i2}, \rho_{i1}] = a_{12}, \quad i = 1, 2, \quad (1.1)$$

$$\rho_{2i}^{-1} \rho_{1i} \rho_{2i} = a_{12}^{-1} \rho_{1i} a_{12}, \quad i = 1, 2, \quad (2.1)$$

$$\rho_{2i} \rho_{1i} \rho_{2i}^{-1} = \rho_{1i}^{-1} a_{12} \rho_{1i} a_{12}^{-1} \rho_{1i}, \quad i = 1, 2, \quad (2.2)$$

$$\rho_{1i}^{-1} \rho_{2i} \rho_{1i} = \rho_{2i} a_{12}^{-1} \rho_{2i} a_{12} \rho_{2i}^{-1}, \quad i = 1, 2, \quad (2.3)$$

$$\rho_{1i} \rho_{2i} \rho_{1i}^{-1} = a_{12} \rho_{2i} a_{12}^{-1}, \quad i = 1, 2, \quad (2.4)$$

$$\rho_{21}^{-1} \rho_{12} \rho_{21} = a_{12}^{-1} \rho_{11} a_{12} \rho_{11}^{-1} \rho_{12} a_{12}, \quad (3.1)$$

$$\rho_{21} \rho_{12} \rho_{21}^{-1} = \rho_{11}^{-1} a_{12} \rho_{11} a_{12}^{-1} \rho_{12} \rho_{11}^{-1} a_{12}^{-1} \rho_{11}, \quad (3.2)$$

$$\rho_{11}^{-1} \rho_{22} \rho_{11} = \rho_{22} \rho_{21} a_{12} \rho_{21}^{-1}, \quad (3.3)$$

$$\rho_{11} \rho_{22} \rho_{11}^{-1} = \rho_{22} a_{12}^{-1}, \quad (3.4)$$

$$\rho_{22}^{-1} \rho_{11} \rho_{22} = a_{12}^{-1} \rho_{11}, \quad (4.1)$$

$$\rho_{22} \rho_{11} \rho_{22}^{-1} = \rho_{12}^{-1} a_{12} \rho_{12} \rho_{11}, \quad (4.2)$$

$$\rho_{12}^{-1} \rho_{21} \rho_{12} = \rho_{22} a_{12}^{-1} \rho_{22}^{-1} \rho_{21} a_{12}^{-1} \rho_{22} a_{12} \rho_{22}^{-1}, \quad (4.3)$$

$$\rho_{12} \rho_{21} \rho_{12}^{-1} = a_{12} \rho_{21} \rho_{22}^{-1} a_{12} \rho_{22} a_{12}^{-1}. \quad (4.4)$$

From Theorem 1.4 follows that the group $B_2(T)$ is generated by elements

$$\sigma_1, \rho_{11}, \rho_{12},$$

which are connected with generators of group $P_2(T)$ by equations

$$a_{12} = \sigma_1^2, \rho_{21} = \sigma_1^{-1} \rho_{11} \sigma_1, \rho_{22} = \sigma_1^{-1} \rho_{12} \sigma_1.$$

A group $A_1 = \langle a_{12}, \rho_{11}, \rho_{12} \rangle$ is isomorphic to a free group with free generators ρ_{11}, ρ_{12} , which is normal in the group $P_2(T)$.

From the relation (1.1) follows that in the group A_1 realizes the relation

$$[\rho_{12}, \rho_{11}] = a_{12}. \quad (5)$$

There conjugate both hand parts of this relation by element ρ_{22} . Then in the left hand part using relation (2.1) and (4.1) we get the equation

$$[\rho_{12}, \rho_{11}]^{\rho_{22}} = [a_{12}^{-1} \rho_{12} a_{12}, a_{12}^{-1} \rho_{11}] = a_{12}^{-1} \rho_{12}^{-1} a_{12} \rho_{11}^{-1} \rho_{12} \rho_{11}.$$

In the right hand part we get the equation

$$\rho_{22}^{-1} a_{12} \rho_{22} = a_{12}^{-1} \rho_{12} a_{12} \rho_{12}^{-1} a_{12}. \quad (6)$$

Really, from the relation (2.4) we have the equation

$$\rho_{22}^{-1} a_{12} \rho_{22} = (\rho_{22}^{-1} \rho_{12} \rho_{22}) \rho_{12}^{-1} a_{12}.$$

Applying to the expression in the brackets relation (2.1) unde $i = 2$ we get the equation (6). Therefore, in the free group A_1 we have the equation

$$a_{12}^{-1} \rho_{12}^{-1} a_{12} \rho_{11}^{-1} \rho_{12} \rho_{11} = a_{12}^{-1} \rho_{12} a_{12} \rho_{12}^{-1} a_{12},$$

or, after multiplication on the left on a_{12} :

$$\rho_{12}^{-1} a_{12} \rho_{11}^{-1} \rho_{12} \rho_{11} = \rho_{12} a_{12} \rho_{12}^{-1} a_{12}. \quad (7)$$

The group A_1 is free generated by elements ρ_{11}, ρ_{12} , and element a_{12} is expressed through these elements with help of equation (5). There escape in the equation (7) from the generator a_{12} we get the equation

$$\rho_{12}^{-2} \rho_{11}^{-1} \rho_{12} = \rho_{11}^{-1} \rho_{12} \rho_{11} \rho_{12}^{-2} \rho_{11}^{-1}.$$

But in the free group with free generators ρ_{11} and ρ_{12} such equation impossible. Hence, the system of defining relations of Scott for the braid groups of two dimension orientable manifolds is contradiction. Similar, there can show that although the system of defining relations which is constructed in Theorems 1.1 1.2 for the braid groups of non-orientable manifolds is contradiction too.

§ 3. Braid group of genetic code

Let us consider a genetic code

$$\mathcal{G} = \langle X \parallel R \rangle, \quad (1)$$

where $X = \{x_1, \dots, x_l\}$ is a finite set of letters and R is an empty set or a set which contains one word in the alphabet $X^{\pm 1} = \{x_1^{\pm 1}, \dots, x_l^{\pm 1}\}$. At that we will distinguish cases when $R = \emptyset$ is an empty set and $R = \{e\}$ is a set from the trivial word. The group which is defined by the genetic code (1) we will denote by $G(\mathcal{G})$. We compare to the genetic code (1) and a natural number n a group $B_n(\mathcal{G})$ which will call *n-strand braid group of genetic code* $\mathcal{G}$. If $n = 1$ then we let $B_1(\mathcal{G}) = G(\mathcal{G})$. If $n > 1$, $R = \{r\}$ then the group $B_n(\mathcal{G})$ is defined by generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$, $x_1, \dots, x_l$ and defining relations

$$\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \quad i = 1, 2, \dots, n-2, \quad (2)$$

$$\sigma_i \sigma_j = \sigma_j \sigma_i, \quad |i - j| \geq 2, \quad (3)$$

$$\sigma_i x_j = x_j \sigma_i, \quad i = 2, 3, \dots, n-1, \quad j = 1, 2, \dots, l, \quad (4)$$

$$x_j (\sigma_1^{-1} x_j \sigma_1^{-1}) = (\sigma_1^{-1} x_j \sigma_1^{-1}) x_j, \quad j = 1, 2, \dots, l, \quad (5)$$

$$x_j (\sigma_1^{-1} x_i^{-1} \sigma_1) x_j^{-1} (\sigma_1^{-1} x_i \sigma_1) = \sigma_1^2, \quad 1 \leq i < j \leq l, \quad (6)$$

$$r = \sigma_1 \sigma_2 \dots \sigma_{n-2} \sigma_{n-1}^2 \sigma_{n-2} \dots \sigma_2 \sigma_1. \quad (7)$$

Such defined class groups $B_n(\mathcal{G})$ includes many known groups. Let us give a few examples.

If $\mathcal{G}_\emptyset = \langle \emptyset \parallel \emptyset \rangle$ is the empty genetic code then there are not exist the relations (4)–(7) and $B_n(\mathcal{G}_\emptyset) = B_n(E^2)$ is the braid group of Euclidean plane. If $\mathcal{G}_0^\emptyset = \langle \emptyset \parallel e = e \rangle$, then there are not exist the relations (4)–(6) and $B_n(\mathcal{G}_0^\emptyset) = B_n(S^2)$ is the braid group of two dimension sphere.

Consider the genetic codes

$$\mathcal{G}_\emptyset^l = \langle x_1, x_2, \dots, x_l \parallel \emptyset \rangle, \quad \mathcal{G}_0^l = \langle x_1, x_2, \dots, x_l \parallel e = e \rangle,$$

$$\mathcal{G}_1^l = \langle x_1, x_2, \dots, x_l \parallel r = e \rangle, \quad l \geq 1,$$

where r is a non-trivial word. At that we consider that the group $G(\mathcal{G}_1^l)$ is not a free group since in the opposite case we can define (IT) by the genetic code with the empty set of relations. Evidently, $G(\mathcal{G}_\emptyset^l) \simeq G(\mathcal{G}_0^l) \simeq F_l$ is the free group with free generators $x_1, x_2, \dots, x_l$; $G(\mathcal{G}_1^l)$ is the group with one defining relation, and this group is a homomorphic image of groups $G(\mathcal{G}_\emptyset^l)$ and $G(\mathcal{G}_0^l)$.

If

$$\mathcal{G}_1^{2p} = \langle x_1, \dots, x_{2p} \parallel x_1 x_2^{-1} x_3 \dots x_{2p}^{-1} x_1^{-1} x_2 x_3^{-1} \dots x_{2p} \rangle, \quad p \geq 1,$$

is the genetic code of fundamental group of compact orientable surface genus p then $B_n(\mathcal{G}_1^{2p})$ is the braid group of this surface, it was defined by Zariski [16]. If $\mathcal{G} = \langle x \parallel \emptyset \rangle$ then $B_n(\mathcal{G})$ is an Artin braid group corresponding to the Coxeter group of type B [31]. Indeed, Coxeter group of type B corresponds to the Artin group with generators

$\sigma_1, \sigma_2, \dots, \sigma_{n-1}, \tau$, where the generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ satisfy to relations (2)–(3), and the relations which contain τ have the form

$$\sigma_i \tau = \tau \sigma_i, \quad i = 2, 3, \dots, n-1, \quad (8)$$

$$\tau \sigma_1 \tau \sigma_1 = \sigma_1 \tau \sigma_1 \tau, \quad (9)$$

i. e. relations (8) correspond to relations (4), and rewriting the relations (9) in the form

$$\tau^{-1} \sigma_1^{-1} \tau^{-1} \sigma_1^{-1} = \sigma_1^{-1} \tau^{-1} \sigma_1^{-1} \tau^{-1},$$

and put $\tau^{-1} = x$ we get the braid group $B_n(\langle x \parallel \emptyset \rangle)$. Note that this group connects to links in body torus similar as the braid group of Euclidean plane connects to links in the three dimension sphere S^3 (see [7]).

The braid group $B_n(\mathcal{G})$ in general case depends from the genetic code $\mathcal{G}$ but not only from the group $G(\mathcal{G})$. For example, $\mathcal{G}_\emptyset^l$ and $\mathcal{G}_0^l$ define the isomorphic groups but the corresponding braid groups $B_n(\mathcal{G}_\emptyset^l)$ and $B_n(\mathcal{G}_0^l)$ are not isomorphic for $n > 1$.

Let's try to clarify how the corresponding braid groups $B_n(\mathcal{G}_\emptyset^l)$, $B_n(\mathcal{G}_0^l)$ and $B_n(\mathcal{G}_1^l)$ are connected for $n \geq 2$.

By the symbol $\mathcal{G}_\emptyset$ we will further denote the genetic code $\mathcal{G}_\emptyset^\emptyset$, or genetic code $\mathcal{G}_\emptyset^l$. By the symbol $\mathcal{G}_0$ we will further denote the genetic code $\mathcal{G}_0^\emptyset$, or genetic code $\mathcal{G}_0^l$. By the symbol $\mathcal{G}_1$ we will denote the genetic code $\mathcal{G}_1^l$, containing a nontrivial relation. To indicate the explicit form of the relation r of the genetic code $\mathcal{G}_1^l$ we will write $\mathcal{G}_1^l(r)$.

Note that in the group $B_n(\mathcal{G}_\emptyset)$ does not hold the relation (7). Hence, there exists homomorphisms $\mu_1 : B_n(\mathcal{G}_\emptyset) \rightarrow B_n(\mathcal{G}_0)$, $\mu_2 : B_n(\mathcal{G}_\emptyset) \rightarrow B_n(\mathcal{G}_1)$. And the kernel of the first homomorphism $\ker(\mu_1)$ is the normal closure of the element

$$\sigma_1 \sigma_2 \dots \sigma_{n-2} \sigma_{n-1}^2 \sigma_{n-2} \dots \sigma_2 \sigma_1$$

in the group $B_n(\mathcal{G}_\emptyset)$, while the kernel of the second homomorphism is the normal closure of the element

$$r^{-1} \sigma_1 \sigma_2 \dots \sigma_{n-2} \sigma_{n-1}^2 \sigma_{n-2} \dots \sigma_2 \sigma_1.$$

For the group $B_n(\mathcal{G})$, $n > 1$, there is a homomorphism ν onto the symmetric group S_n which is defined on the generators by equations:

$$\nu(\sigma_i) = (i, i+1), \quad i = 1, 2, \dots, n-1; \quad \nu(x_j) = 1, \quad j = 1, 2, \dots, l.$$

The kernel of this homomorphism $\ker(\nu)$ we will call *the pure braid group* on n strands and denote by $P_n(\mathcal{G})$.

To define the structure of $B_n(\mathcal{G})$ we define its subgroup $D_n = D_n(\mathcal{G})$ which includes such elements that under the action of homomorphism ν map to permutations which fixing the symbol 1.

Let us introduce the following definitions:

$$x_{1i} = x_i, \quad x_{k+1,i} = \sigma_k^{-1} x_{ki} \sigma_k, \quad i = 1, 2, \dots, l, \quad 1 \leq k \leq n-1.$$

In this definitions the following lemma is true, and it is an analog of the corresponding lemma from the paper of Zariski.

Lemma 1. *In the group D_n the following formulas of conjugation are true*

- 1) $\sigma_k^\varepsilon x_{1i} \sigma_k^{-\varepsilon} = x_{1i}, \quad \varepsilon = \pm 1, \quad k = 2, 3, \dots, n-1; \quad i = 1, 2, \dots, l$
- 2) $x_{2i} a_{12} x_{2i}^{-1} = x_{1i}^{-1} a_{12} x_{1i},$
- 3) $x_{2i}^{-1} a_{12} x_{2i} = a_{12}^{-1} x_{1i} a_{12} x_{1i}^{-1} a_{12},$
- 4) $x_{2i}^{-\varepsilon} a_{1j} x_{2i}^\varepsilon = a_{1j}, \quad j > 2,$
- 5) $x_{2i}^{-1} x_{1i} x_{2i} = a_{12}^{-1} x_{1i} a_{12},$
- 6) $x_{2i} x_{1i} x_{2i}^{-1} = x_{1i}^{-1} a_{12} x_{1i} a_{12}^{-1} x_{1i},$
- 7) $x_{2i}^{-1} x_{1j} x_{2i} = a_{12}^{-1} x_{1j}, \quad i < j,$
- 8) $x_{2i} x_{1j} x_{2i}^{-1} = (x_{1i}^{-1} a_{12} x_{1i}) x_{1j}, \quad i < j,$
- 9) $x_{2j} x_{1i} x_{2j}^{-1} = (x_{1j}^{-1} a_{12} x_{1j}) a_{12}^{-1} x_{1i} (x_{1j}^{-1} a_{12}^{-1} x_{1j}), \quad i < j,$
- 10) $x_{2j}^{-1} x_{1i} x_{2j} = (a_{12}^{-1} x_{1j} a_{12} x_{1j}^{-1}) x_{1i} a_{12}, \quad i < j.$

Proof. The formula 1) follows from the defining relation (4).

From the relation (5) follows the relation

$$x_{1i} (\sigma_1^{-1} x_{1i} \sigma_1^{-1}) = (\sigma_1^{-1} x_{1i} \sigma_1^{-1}) x_{1i}, \quad i = 1, 2, \dots, l. \quad (10)$$

From this

$$(\sigma_1 x_{1i}^{-1} \sigma_1) x_{1i} (\sigma_1^{-1} x_{1i} \sigma_1^{-1}) = x_{1i}, \quad i = 1, 2, \dots, l.$$

Conjugating both hand sides of this equation by element $a_{12} = \sigma_1^2$ we get the equation 5). Then conjugating the equation 5) by element x_{2i}^{-1} we get

$$x_{1i} = x_{2i} a_{12}^{-1} x_{1i} a_{12} x_{2i}^{-1}. \quad (11)$$

On the other hand from (10) follows that

$$\sigma_1^{-1} x_{1i} \sigma_1^{-1} x_{1i} = a_{12}^{-1} x_{1i} \sigma_1^{-1} x_{1i} \sigma_1,$$

i. e.

$$(\sigma_1^{-1} x_{1i} \sigma_1) a_{12}^{-1} x_{1i} = a_{12}^{-1} x_{1i} (\sigma_1^{-1} x_{1i} \sigma_1).$$

Since $\sigma_1^{-1} x_{1i} \sigma_1 = x_{2i}$ we see that the elements x_{2i} and $a_{12}^{-1} x_{1i}$ are permutable. Hence, we can write (11) in the form

$$x_{1i} = a_{12}^{-1} x_{1i} x_{2i} a_{12} x_{2i}^{-1},$$

i. e.

$$x_{1i}^{-1} a_{12} x_{1i} = x_{2i} a_{12} x_{2i}^{-1}, \quad (12)$$

but this is the equation 2).

If we conjugate both hand sides of relation (12) by element σ_1 then we get

$$(\sigma_1^{-1} x_{1i}^{-1} \sigma_1) a_{12} (\sigma_1^{-1} x_{1i} \sigma_1) = a_{12}^{-1} x_{1i} a_{12} x_{1i}^{-1} a_{12}$$

and the equation 3) is ascertained.

To prove 4) we remember that $a_{1j} = \sigma_1^{-1} \dots \sigma_{j-2}^{-1} \sigma_{j-1}^2 \sigma_{j-2} \dots \sigma_1$ and so

$$x_{2i}^\varepsilon a_{1j} x_{2i}^{-\varepsilon} = (\sigma_1^{-1} x_{1i}^\varepsilon \sigma_1) a_{1j} (\sigma_1^{-1} x_{1i}^{-\varepsilon} \sigma_1) = \sigma_1^{-1} x_{1i}^\varepsilon a_{2j} x_{1i}^{-\varepsilon} \sigma_1 = \sigma_1^{-1} a_{2j} \sigma_1 = a_{1j},$$

i. e. the equation 4) is true.

Conjugate both hand sides of the equation 5) by element $a_{12}^{-1} x_{1i}$, we get

$$x_{1i}^{-1} a_{12} x_{2i}^{-1} x_{1i} x_{2i} a_{12}^{-1} x_{1i} = x_{1i}.$$

Using that element x_{2i} is permutable with $a_{12}^{-1} x_{1i}$ we rewrite the last equation in the form

$$x_{2i}^{-1} x_{1i}^{-1} a_{12} x_{1i} a_{12}^{-1} x_{1i} x_{2i} = x_{1i}.$$

Conjugate both hand sides of this relation by element x_{2i}^{-1} we get the relation 6).

Observe that the defining relation (6) of $B_n(\mathcal{G})$ we can write in the form

$$x_{1j} x_{2i}^{-1} x_{1j}^{-1} x_{2i} = a_{12}, \quad j > i.$$

From this equation follows 7).

To prove 8) we conjugate both hand sides of equation 7) by element x_{2i}^{-1} , we get

$$x_{1j} = x_{2i} a_{12}^{-1} x_{1j} x_{2i}^{-1}, \quad j > i.$$

Multiplying both hand sides of this equation on the left by $x_{2i} a_{12} x_{2i}^{-1}$ and using 2) we get the required equation.

Conjugating both hand sides of the equation

$$x_{1j} x_{2i}^{-1} x_{1j}^{-1} x_{2i} = a_{12}, \quad j > i,$$

which is true in the group $B_n(\mathcal{G})$ (see relation (6)) by the generator σ_1 , we get

$$x_{2j} a_{12}^{-1} x_{1i}^{-1} a_{12} x_{2j}^{-1} a_{12}^{-1} x_{1i} a_{12} = a_{12}.$$

Multiplying both hand sides of this equation on the right by $a_{12}^{-1} x_{1i}^{-1} a_{12}$ and turning to inverse elements we get the equation

$$(x_{2j} a_{12}^{-1} x_{2j}^{-1})(x_{2j} x_{1i} x_{2j}^{-1})(x_{2j} a_{12} x_{2j}^{-1}) = a_{12}^{-1} x_{1i}.$$

From this

$$x_{2j} x_{1i} x_{2j}^{-1} = (x_{2j} a_{12} x_{2j}^{-1}) a_{12}^{-1} x_{1i} (x_{2j} a_{12}^{-1} x_{2j}^{-1}).$$

Using the equation 2) we get 9).

To prove 10) we conjugate both hand sides of 9) by x_{2j} :

$$\begin{aligned} x_{1i} &= (x_{2j}^{-1} x_{1j}^{-1} x_{2j})(x_{2j}^{-1} a_{12} x_{2j})(x_{2j}^{-1} x_{1j} x_{2j})(x_{2j}^{-1} a_{12}^{-1} x_{2j})(x_{2j}^{-1} x_{1i} x_{2j}) \times \\ &\quad \times (x_{2j}^{-1} x_{1j}^{-1} x_{2j})(x_{2j}^{-1} a_{12}^{-1} x_{2j})(x_{2j}^{-1} x_{1j} x_{2j}). \end{aligned}$$

Using the equations 3) and 5) we get the equation

$$x_{1i} = x_{1j} a_{12}^{-1} x_{1j}^{-1} a_{12} (x_{2j}^{-1} x_{1i} x_{2j}) a_{12}^{-1}.$$

From this

$$x_{2j}^{-1} x_{1i} x_{2j} = (a_{12}^{-1} x_{1j} a_{12} x_{1j}^{-1}) x_{1i} a_{12},$$

but it is the equation 10). The lemma is proven.

Lemma 2. 1) As a set of right coset representatives of D_n in $B_n(\mathcal{G})$ we can take the set $M_n = \{1, \sigma_1, \sigma_2 \sigma_1, \dots, \sigma_{n-1} \sigma_{n-2} \dots \sigma_2 \sigma_1\}$. 2) The group D_n is generated by elements $a_{12}, a_{13}, \dots, a_{1n}, \sigma_2, \sigma_3, \dots, \sigma_{n-1}, x_{1k}, x_{2k}, k = 1, 2, \dots, l$. 3) The subgroup $U_n = \langle a_{12}, a_{13}, \dots, a_{1n}, x_{11}, x_{12}, \dots, x_{1l} \rangle$ is normal in D_n .

Proof. 1) We denote by $m_0 = 1$, $m_i = \sigma_i \sigma_{i-1} \dots \sigma_2 \sigma_1$, $i = 1, 2, \dots, n-1$. Note that the image of the element m_i in S_n maps the symbol 1 to $i+1$ (we view the multiplication of permutations from right to left). So M_n is a set of right coset representatives of D_n in $B_n(\mathcal{G})$. And what is more, M_n is a Schreier set of coset representatives.

To prove 2) we use the Reidemeister-Schreier method. Let $u \mapsto \bar{u}$ be a function which takes the right coset representatives of D_n in $B_n(\mathcal{G})$. Then

$$D_n = \langle d_{x,m} = (\overline{xm})^{-1}xm \mid m \in M_n, x \in \{\sigma_1, \sigma_2, \dots, \sigma_{n-1}, x_1, x_2, \dots, x_l\} \rangle.$$

We will show that if $x = \sigma_i$ for some $i = 1, 2, \dots, n-1$ then

$$d_{\sigma_i, m_j} = \begin{cases} \sigma_i & \text{if } i > j+1, \\ 1 & \text{if } i = j+1, \\ a_{1,i+1} & \text{if } i = j, \\ \sigma_{i+1} & \text{if } i < j. \end{cases}$$

Indeed, for $i > j+1$ we have

$$d_{\sigma_i, m_j} = (\overline{\sigma_i m_j})^{-1} \sigma_i m_j = m_j^{-1} \sigma_i m_j = \sigma_i.$$

For $i = j+1$

$$d_{\sigma_i, m_{i-1}} = (\overline{\sigma_i m_{i-1}})^{-1} \sigma_i m_{i-1} = m_{i-1}^{-1} \sigma_i m_{i-1} = m_i^{-1} m_i = 1.$$

For $i = j$

$$d_{\sigma_i, m_i} = (\overline{\sigma_i \sigma_i m_{i-1}})^{-1} \sigma_i^2 m_{i-1} = m_{i-1}^{-1} \sigma_i^2 m_{i-1} = m_{i-2}^{-1} \sigma_i \sigma_{i-1}^2 \sigma_i^{-1} m_{i-2} =$$

$$= \sigma_i m_{i-3}^{-1} (\sigma_{i-2}^{-1} \sigma_{i-1}^2 \sigma_{i-2}) m_{i-3} \sigma_i^{-1} = \sigma_i m_{i-3}^{-1} \sigma_{i-1} \sigma_{i-2}^2 \sigma_{i-1}^{-1} m_{i-3} \sigma_i^{-1} =$$

$$\sigma_i \sigma_{i-1} m_{i-3}^{-1} \sigma_{i-2}^2 m_{i-3} \sigma_{i-1}^{-1} \sigma_i = \dots = \sigma_i \sigma_{i-1} \dots \sigma_2 \sigma_1^2 \sigma_2^{-1} \dots \sigma_{i-1}^{-1} \sigma_i^{-1} = a_{1i}.$$

At the end, for $i < j$ the requirement equation is got from the following line:

$$\begin{aligned} d_{\sigma_i, m_j} &= (\overline{\sigma_i m_j})^{-1} \sigma_i m_j = m_j^{-1} \sigma_i m_j = \\ &= \sigma_1^{-1} \sigma_2^{-1} \dots \sigma_{i+1}^{-1} (\sigma_{i+2}^{-1} \dots \sigma_j^{-1} \sigma_i \sigma_j \dots \sigma_{i+2}) \sigma_{i+1} \dots \sigma_2 \sigma_1 = \\ &= m_{i+1}^{-1} \sigma_i m_{i+1} = m_i^{-1} \sigma_{i+1}^{-1} (\sigma_i \sigma_{i+1} \sigma_i) m_{i-1} = \\ &= m_i^{-1} \sigma_{i+1}^{-1} (\sigma_i \sigma_{i+1} \sigma_i) m_{i-1} = m_i^{-1} \sigma_{i+1}^{-1} \sigma_{i+1} \sigma_i \sigma_{i+1} m_{i-1} = \\ &= m_i^{-1} \sigma_i m_{i-1} \sigma_{i+1} = m_i^{-1} m_i \sigma_{i+1} = \sigma_{i+1}. \end{aligned}$$

Further, if $x = x_i$, $i = 1, 2, \dots, l$, then

$$d_{x_i, m_j} = \begin{cases} x_i & \text{if } j = 0, \\ x_{2i} & \text{if } j = 1, 2, \dots, n-1. \end{cases}$$

Indeed,

$$\begin{aligned} d_{x_i, 1} &= (\overline{x_i})^{-1} x_i = x_i, \quad d_{x_i, m_j} = \overline{m_j}^{-1} x_i m_j = \sigma_1^{-1} x_i \sigma_1 = x_{2i}, \\ &\quad i = 1, 2, \dots, l, \quad j = 1, 2, \dots, n-1. \end{aligned}$$

Hence, the item 2) of lemma is proven.

The item 3) immediately follows from the item 2), conjugation rules in B_n , and lemma 1.

Using this lemma we can represent every element $w \in B_n(\mathcal{G})$ as the product

$$w = w_1 m_{n,i_1}, \quad w_1 \in D_n, \quad m_{n,i_1} = m_{i_1} = \sigma_{i_1} \sigma_{i_1-1} \dots \sigma_1 \in M_n,$$

and the element w_1 in one's part we can represent in the form

$$w_1 = w_2 u_n,$$

where $w_2 \in \langle \sigma_2, \sigma_3, \dots, \sigma_{n-1}, x_{21}, x_{22}, \dots, x_{2l} \rangle$, $u_n \in U_n$. Denote by $\overline{B}_{n-1}$ a subgroup of $B_n(\mathcal{G})$ which is generated by elements $\sigma_2, \sigma_3, \dots, \sigma_{n-1}, x_{21}, x_{22}, \dots, x_{2l}$. It is true

Lemma 3. *In the group $\overline{B}_{n-1} = \langle \sigma_2, \sigma_3, \dots, \sigma_{n-1}, x_{21}, x_{22}, \dots, x_{2l} \rangle$ the following relations are true*

$$\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \quad i = 2, 3, \dots, n-2, \quad (13)$$

$$\sigma_i \sigma_j = \sigma_j \sigma_i, \quad |i - j| \geq 2, \quad (14)$$

$$\sigma_i x_{2k} = x_{2k} \sigma_i, \quad i = 3, 4, \dots, n-1, \quad k = 1, 2, \dots, l, \quad (15)$$

$$x_{2k} (\sigma_2^{-1} x_{2k} \sigma_2^{-1}) = (\sigma_2^{-1} x_{2k} \sigma_2^{-1}) x_{2k}, \quad k = 1, 2, \dots, l, \quad (16)$$

$$x_{2j} (\sigma_2^{-1} x_{2i}^{-1} \sigma_2) x_{2j}^{-1} (\sigma_2^{-1} x_{2i} \sigma_2) = \sigma_2^2, \quad 1 \leq i < j \leq l, \quad (17)$$

$$r_2 = \sigma_2 \sigma_3 \dots \sigma_{n-2} \sigma_{n-1}^2 \sigma_{n-2} \dots \sigma_3 \sigma_2 a_{12}. \quad (18)$$

where the word r_2 is got from $r_1 = r$ conjugation by element σ_1 , i. e. if we replace all elements $x_{1k} = x_k$, which be included in r by elements x_{2k} , $k = 1, 2, \dots, l$.

Proof. Correctness of the relations (13)–(14) is evident. The relation (15) is got from the relation (4) for $i = 3, 4, \dots, n-1$ conjugation by the generator σ_1 . To prove the relation (16) we consider the relation (5):

$$x_{1k} (\sigma_1^{-1} x_{1k} \sigma_1^{-1}) = (\sigma_1^{-1} x_{1k} \sigma_1^{-1}) x_{1k}, \quad k = 1, 2, \dots, l,$$

from $B_n(\mathcal{G})$. Conjugate its both hand sides by element σ_2 and using permutability σ_2 and x_{1k} we get

$$x_{1k} (\sigma_2^{-1} \sigma_1^{-1} \sigma_2 x_{1k} \sigma_2^{-1} \sigma_1^{-1} \sigma_2) = (\sigma_2^{-1} \sigma_1^{-1} \sigma_2 x_{1k} \sigma_2^{-1} \sigma_1^{-1} \sigma_2) x_{1k}.$$

Using the relation $\sigma_2^{-1} \sigma_1^{-1} \sigma_2 = \sigma_1 \sigma_2^{-1} \sigma_1^{-1}$ we get

$$x_{1k} (\sigma_1 \sigma_2^{-1} \sigma_1^{-1} x_{1k} \sigma_1 \sigma_2^{-1} \sigma_1^{-1}) = (\sigma_1 \sigma_2^{-1} \sigma_1^{-1} x_{1k} \sigma_1 \sigma_2^{-1} \sigma_1^{-1}) x_{1k}.$$

Conjugating both hand sides of this equation by element σ_1 we have

$$(\sigma_1^{-1} x_{1k} \sigma_1) \sigma_2^{-1} (\sigma_1^{-1} x_{1k} \sigma_1) \sigma_2^{-1} = \sigma_2^{-1} (\sigma_1^{-1} x_{1k} \sigma_1) \sigma_2^{-1} (\sigma_1^{-1} x_{1k} \sigma_1).$$

Since $\sigma_1^{-1} x_{1k} \sigma_1 = x_{2k}$ then from this we get the relation (16).

To prove the relation (17) we consider the relation

$$x_{1j} (\sigma_1^{-1} x_{1i}^{-1} \sigma_1) x_{1j}^{-1} (\sigma_1^{-1} x_{1i} \sigma_1) = \sigma_1^2, \quad j > i,$$

which is true in $B_n(\mathcal{G})$ (see the relation (6)). Conjugating both hand sides of this relation by element σ_2 , we get

$$\sigma_2^{-1}x_{1j}(\sigma_1^{-1}x_{1i}^{-1}\sigma_1)x_{1j}^{-1}(\sigma_1^{-1}x_{1i}\sigma_1)\sigma_2 = \sigma_2^{-1}\sigma_1^2\sigma_2.$$

Take advantage of permutability σ_2 with every generator x_{1k} we rewrite the last equality in the following form

$$x_{1j}(\sigma_2^{-1}\sigma_1^{-1}\sigma_2)x_{1i}^{-1}(\sigma_2^{-1}\sigma_1\sigma_2)x_{1j}^{-1}(\sigma_2^{-1}\sigma_1^{-1}\sigma_2)x_{1i}(\sigma_2^{-1}\sigma_1\sigma_2) = \sigma_2^{-1}\sigma_1^2\sigma_2.$$

Using the equations

$$\sigma_2^{-1}\sigma_1^{-1}\sigma_2 = \sigma_1\sigma_2^{-1}\sigma_1^{-1}, \quad \sigma_2^{-1}\sigma_1\sigma_2 = \sigma_1\sigma_2\sigma_1^{-1},$$

we get

$$x_{1j}(\sigma_1\sigma_2^{-1}\sigma_1^{-1})x_{1i}^{-1}(\sigma_1\sigma_2\sigma_1^{-1})x_{1j}^{-1}(\sigma_1\sigma_2^{-1}\sigma_1^{-1})x_{1i}(\sigma_1\sigma_2\sigma_1^{-1}) = \sigma_1\sigma_2^2\sigma_1^{-1}.$$

Conjugating both hand sides of this equation by element σ_1 , and using that $x_{2k} = \sigma_1^{-1}x_{1k}\sigma_1$, $k = 1, 2, \dots, l$, we get the relation (17).

The relation (18) is obtained from the relation (7) conjugation by element σ_1 . The lemma is proven.

From this lemma follows that the group $\overline{B}_{n-1}$ is isomorphic to the braid group $B_{n-1}(\mathcal{G})$ in the case then there is not the relation (7). Hence, in this case the following assertion is true

Lemma 4. *For the genetic code $\mathcal{G} = \mathcal{G}_\emptyset$ (doesn't contain relations) the group D_n contains a subgroup which is isomorphic to $B_{n-1}(\mathcal{G})$ and so D_n is the semi-direct product: $D_n = U_n \rtimes B_{n-1}(\mathcal{G})$.*

In the case when $\mathcal{G} = \mathcal{G}_1$ (contains only one non-trivial relation) it is easy to see that the factor-group D_n/U_n is isomorphic to $B_{n-1}(\mathcal{G})$, but in this case we can't assert that $B_{n-1}(\mathcal{G})$ is a subgroup of the group $B_n(\mathcal{G})$.

Further we can consider the group $B_{n-1}(\mathcal{G}) = \langle \sigma_2, \dots, \sigma_{n-1}, x_{21}, \dots, x_{2l} \rangle$, its homomorphism onto the symmetric group S_n which is induced by the homomorphism $\nu : B_n(\mathcal{G}) \longrightarrow S_n$, choose the subgroup $D_{n-1} \leq B_{n-1}(\mathcal{G})$ that consists from the elements, which under the action of ν map to the permutations that fix the symbol 2. As a set of coset representatives of D_{n-1} in $B_{n-1}(\mathcal{G})$ we can take the set

$$M_{n-1} = \{1, \sigma_2, \sigma_3\sigma_2, \dots, \sigma_{n-1}\sigma_{n-2} \dots \sigma_2\}.$$

As above, we can show that D_{n-1} is generated by elements

$$a_{23}, a_{24}, \dots, a_{2n}, \sigma_3, \sigma_4, \dots, \sigma_{n-1}, x_{2k}, x_{3k}, \quad k = 1, 2, \dots, l,$$

and the subgroup $U_{n-1} = \langle a_{23}, a_{24}, \dots, a_{2n}, x_{21}, x_{22}, \dots, x_{2l} \rangle$ is normal in D_{n-1} . In this case the factor-group D_{n-1}/U_{n-1} is isomorphic to $B_{n-2}(\mathcal{G})$. Continuing this process, on the last step we get a group $B_1(\mathcal{G})$ which is isomorphic to $G(\mathcal{G})$. In that way, for any $i = 0, 1, \dots, n-2$ we construct a sequence

$$B_{n-i}(\mathcal{G}) \geq D_{n-i} \supseteq U_{n-i},$$

where

$$\begin{aligned} B_{n-i}(\mathcal{G}) &= \langle \sigma_{i+1}, \sigma_{i+2}, \dots, \sigma_{n-1}, x_{i+1,1}, \dots, x_{i+1,l} \rangle, \\ D_{n-i} &= \langle a_{i+1,i+2}, a_{i+1,i+3}, \dots, a_{i+1,n}, \sigma_{i+2}, \sigma_{i+3}, \dots, \sigma_{n-1}, x_{i+1,1}, \dots, \\ &\quad x_{i+1,l}, x_{i+2,1}, \dots, x_{i+2,l} \rangle, \\ U_{n-i} &= \langle a_{i+1,i+2}, a_{i+1,i+3}, \dots, a_{i+1,n}, x_{i+1,1}, x_{i+1,2}, \dots, x_{i+1,l} \rangle. \end{aligned}$$

§ 4. Some properties of $B_n(\mathcal{G})$

We will show that $B_n(\mathcal{G}_0)$ for $n \geq 2$ has a torsion. It is true

Theorem 1. *The group $B_n(\mathcal{G}_0)$ has a torsion for all $n \geq 2$. The subgroup $U_3(\mathcal{G}_0)$ has elements of order 2. The subgroups $U_n(\mathcal{G}_0)$ for $n \neq 3$ are free.*

Proof. Indeed, for $n = 2$ in the group $B_n(\mathcal{G}_0)$ the following relation $a_{12} = 1$ is true, i. e. $\sigma_1^2 = 1$. If $\mathcal{G}_0 = \mathcal{G}_0^\emptyset$ then $B_2(\mathcal{G}_0) \simeq \mathbb{Z}_2$. If $\mathcal{G}_0 = \mathcal{G}_0^l$ then the group $B_2(\mathcal{G}_0)$ is generated by elements $x_1, x_2, \dots, x_l, \sigma_1$. Let $x_{1k} = x_k$, $x_{2k} = \sigma_1^{-1}x_{1k}\sigma_1 = \sigma_1x_{1k}\sigma_1$, $k = 1, 2, \dots, l$. Then relations (5)–(7) have the following form:

$$x_{1j}x_{2j} = x_{2j}x_{1j}, \quad j = 1, 2, \dots, l,$$

$$x_{1j}x_{2i}^{-1} = x_{2i}^{-1}x_{1j}, \quad j > i,$$

$$\sigma_1^2 = 1.$$

From these relations follows that the pure braid group $P_2(\mathcal{G}_0^l) \simeq F_l \times F_l$ is the direct product of two copies of the free group F_l . The group U_2 is generated by elements x_{1k} , $k = 1, 2, \dots, l$, and so is free, and $U_1 = \langle x_{21}, x_{22}, \dots, x_{2l} \rangle$ is free too.

By the lemma 6 in $B_n(\mathcal{G}_0)$ for $n = 3$ the following relations

$$a_{12}a_{13} = 1, \quad a_{23}a_{12} = 1, \quad a_{13}a_{23} = 1$$

are true. If we express from the first relation a_{12} and substitute it in the second then we get $a_{23}a_{13}^{-1} = 1$, i. e. $a_{23} = a_{13}$. Then from the third relation we have $a_{23}^2 = 1$. Hence, $a_{13} = a_{23}$, $a_{12} = a_{13}^{-1} = a_{23}^{-1} = a_{23}$. Then

$$U_3 = \langle a_{13}, x_{11}, x_{12}, \dots, x_{1l} \rangle \simeq \mathbb{Z}_2 * F_l,$$

$$U_2 = \langle x_{21}, x_{22}, \dots, x_{2l} \rangle \simeq F_l,$$

$$U_1 = \langle x_{31}, x_{32}, \dots, x_{3l} \rangle \simeq F_l \simeq G(\mathcal{G}_0).$$

For $n = 4$ in the group $B_n(\mathcal{G}_0)$ the following relations hold

$$a_{12}a_{13}a_{14} = 1, \quad a_{23}a_{24}a_{12} = 1, \quad a_{34}a_{13}a_{23} = 1, \quad a_{14}a_{24}a_{34} = 1.$$

Let us express from the first relation a_{14} , from the second a_{24} , from the third a_{34} and substitute these expressions in the last relation, we get

$$a_{14}a_{24}a_{34} = (a_{12}^{-1}a_{23}^{-1}a_{13}^{-1})^2 = 1,$$

i. e.

$$(a_{13}a_{23}a_{12})^2 = (a_{12}a_{13}a_{23})^2 = 1.$$

Since $a_{12}a_{13}a_{23} = (\sigma_1\sigma_2)^3$, and the image of the element $\sigma_1\sigma_2$ in the group S_4 different from 1 then the group $B_4(\mathcal{G}_0)$ contains some elements of finite order.

Consider the subgroup

$$U_4 = \langle a_{12}, a_{13}, a_{14}, x_{11}, x_{12}, \dots, x_{1l} \rangle.$$

In this group the relation $a_{12}a_{13}a_{14} = 1$ is true. If we exclude from it a_{12} , then we see that

$$U_4 = \langle a_{13}, a_{14}, x_{11}, x_{12}, \dots, x_{1l} \rangle$$

is a free group. Further,

$$U_3 = \langle a_{23}, a_{24}, x_{21}, x_{22}, \dots, x_{2l} \mid a_{23}a_{24} = 1 \rangle = \langle a_{24}, x_{21}, x_{22}, \dots, x_{2l} \rangle,$$

but since $(a_{12}a_{13}a_{23})^2 = 1$ then in the group U_3 the following equation $a_{23}^2 = a_{24}^{-2} = 1$ is true. Hence, $U_3 \simeq \mathbb{Z}_2 * F_l$.

Consider

$$U_2 = \langle a_{34}, x_{31}, x_{32}, \dots, x_{3l} \mid a_{34} = 1 \rangle = \langle x_{31}, x_{32}, \dots, x_{3l} \rangle.$$

Hence, U_2 as well as the group $U_1 = G(\mathcal{G}_0)$ is free.

Further, for arbitrary $n \geq 4$ in view of the lemma 6 the following equations

$$a_{12}a_{13} \dots a_{1n} = 1,$$

$$a_{23}a_{24} \dots a_{2n}a_{12} = 1,$$

$$a_{34}a_{35} \dots a_{3n}a_{13}a_{23} = 1,$$

$$\dots\dots\dots$$

$$a_{n-1,n}a_{1,n-1}a_{2,n-1} \dots a_{n-2,n-1} = 1,$$

$$a_{1n}a_{2n} \dots a_{n-1,n} = 1,$$

are true in the group $B_n(\mathcal{G}_0)$. Express from the first equation a_{1n} , from the second a_{2n} , from the third a_{3n} , and so on, finally, from $n-1$ -th we express $a_{n-1,n}$. If we substitute these expressions in the last equation and use the defining relations of the group P_n then we get

$$[a_{12}^{-1}(a_{23}^{-1}a_{13}^{-1})(a_{34}^{-1}a_{24}^{-1}a_{14}^{-1}) \dots (a_{n-2,n-1}^{-1}a_{n-3,n-1}^{-1} \dots a_{1,n-1}^{-1})]^2 = 1.$$

It is not hard to check that an element which is inverse to the element from the square bracket, is equal to

$$a_{12}(a_{13}a_{23})(a_{14}a_{24}a_{34}) \dots (a_{1,n-1}a_{2,n-1} \dots a_{n-2,n-1}),$$

which is equal to

$$(\sigma_1\sigma_2 \dots \sigma_{n-2})^{n-1}$$

(see [25, c. 28]). Hence, $(\sigma_1\sigma_2 \dots \sigma_{n-2})^{2n-2} = 1$. It is easy to see that the image of the element $\sigma_1\sigma_2 \dots \sigma_{n-2}$ in the group S_n different from 1, and so the group $B_n(\mathcal{G}_0)$ has elements of finite order.

Now consider the subgroups U_{n-i+1} , which have the genetic code

$$U_{n-i+1} = \langle a_{i,i+1}, a_{i,i+2}, \dots, a_{i,n}x_{i1}, x_{i2}, \dots, x_{il} \mid a_{i,i+1}a_{i,i+2} \dots a_{i,n} = 1 \rangle.$$

If we exclude from the relation the generator $a_{i,i+1}$ then we see that

$$U_{n-i+1} = \langle a_{i,i+2}, a_{i,i+3}, \dots, a_{i,n}, x_{i1}, x_{i2}, \dots, x_{il} \rangle, \quad i = 1, 2, \dots, n-1.$$

For all $i \neq n-2$ this group is free, and for $i = n-2$ we have

$$U_3 = \langle a_{n-2,n-1}, a_{n-2,n}, x_{n-2,1}, x_{n-2,2}, \dots, x_{n-2,l} \mid a_{n-2,n-1}a_{n-2,n} = 1, a_{n-2}^2 = 1 \rangle \simeq \mathbb{Z}_2 * F_l.$$

Thus the theorem is proven.

This theorem generalizes a result from [18] which asserts that the braid group of sphere $B_n(S^2)$ has a torsion.

If we use the same argumentation as in proof of the theorem 1 then we easy get

Corollary. *The subgroup $U_n(\mathcal{G}_1)$ of the braid group $B_n(\mathcal{G}_1)$, $n \geq 2$, is free.*

Similar to the classical braid groups it is holds

Theorem 2. *In the group $B_n(\mathcal{G})$ the word problem is solable.*

Proof. Let a word w be written in the generators of $B_n(\mathcal{G})$. As was proved in the previous section there exists the short exact sequence

$$1 \longrightarrow P_n(\mathcal{G}) \longrightarrow B_n(\mathcal{G}) \longrightarrow S_n \longrightarrow 1.$$

If the image of w in S_n is non-trivial then $w \neq 1$. In the opposite case, $w \in P_n(\mathcal{G})$. Let us rewrite this word in the generators of the group $P_n(\mathcal{G})$. Then we represent w as a product $w = w_1 w'$, where $w_1 \in U_n$ and is a word in generators a_{1j} , $j = 2, 3, \dots, n$, x_{1k} , $k = 1, 2, \dots, l$, and the word w' doesn't contain these generators. Now we consider the short exact sequence

$$1 \longrightarrow U_n \longrightarrow P_n(\mathcal{G}) \longrightarrow P_{n-1}(\mathcal{G}) \longrightarrow 1.$$

The word $w_1 \in U_n$, and U_n either a free group or is a free product with amalgamation, and hence we can effectively check: is the word w_1 represents trivial element. If $w_1 \neq 1$ then $w \neq 1$. If $w_1 = 1$ then we find an image of element w' in the group $P_{n-1}(\mathcal{G})$, and represent it as a product $w_2 w''$, where w_2 is a word in the generators of U_{n-1} , and w'' doesn't contain these generators. If $w_2 \neq 1$ then $w \neq 1$. If $w_2 = 1$ then we find the image of w'' in the group $P_{n-2}(\mathcal{G})$. If we continue this procedure then we either prove that $w \neq 1$ or on a n -th step we construct an element $w_n \in U_1$. Since $U_1 \simeq G(\mathcal{G})$ and the word problem is effectively solvable in $G(\mathcal{G})$ then we can check: is $w_n = 1$ or not. If $w_n = 1$ then $w = 1$. In the opposite case $w \neq 1$. The theorem is proven.

Recall, that the *rang* $\text{rk}(G)$ of a finitely generated group G [27, p. 29] is called the minimal number of generators of G . For the trivial group G we let $\text{rk}(G) = 0$. It is holds

Proposition 1. *For the braid group $B_n(\mathcal{G})$ the following evaluations hold*

$$\text{rk}(B_1(\mathcal{G})) = \text{rk}(G(\mathcal{G})),$$

$$\text{rk}(B_2(\mathcal{G})) \leq \text{rk}(G(\mathcal{G})) + 1,$$

$$\text{rk}(B_n(\mathcal{G})) \leq \text{rk}(G(\mathcal{G})) + 2, \text{ for } n \geq 2.$$

Proof. The first equality follows from the definition since in this case $B_1(\mathcal{G}) = G(\mathcal{G})$. For $n = 2$ the group $B_2(\mathcal{G})$ is generated by element σ_1 and generators of $G(\mathcal{G})$, and hence the second evaluation holds. The third evaluation follows from the fact that a subgroup of $B_n(\mathcal{G})$ which is generated by elements $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ can be generated by two elements [28, Chapter 6]. The proposition is proven.

In [29] was studied a question about the width of verbal subgroups of the braid group $B_n(E^2)$ of Euclidean plane E^2 , and in [30] similar question was studied for other Artin groups. Recall some facts from these papers.

The *width* $\text{wid}(G, V)$ of verbal subgroup $V(G)$ which is defined in a group G by a set of words V , with respect to this V is called a minimal $m \in \mathbb{N} \cup \{+\infty\}$ such that every element from the subgroup $V(G)$ is a product of $\leq m$ values of words from $V \cup V^{-1}$. We will consider only finite sets of words V since for any verbal subgroup $V(G)$ we can construct some set of words W (which is infinite in general case) such that $W(G) = V(G)$ and $\text{wid}(G, W) = 1$.

We will say that a group G is *rich by verbal subgroups* if every set of words V that define a proper verbal subgroup of a free group F_2 rang two, define a proper verbal subgroup of G (by the proper subgroup we mean a subgroup that is not trivial and is not all group). In particular, if G contains a free non-abelian group then G is rich by verbal subgroups.

We will say that a group G *doesn't have proper verbal subgroups of finite width* if for every finite set V that defines a proper verbal subgroup $V(G)$ of G the width $\text{wid}(G, V)$ is infinite.

Following statements hold

Lemma 7 ([30, Lemma 1]). *Let $G = G_1 \times G_2 \times \dots \times G_k$ be a direct product of finite number subgroups G_i . Then the width $\text{wid}(G, V)$ is equal to the maximal width of the factors $\text{wid}(G_i, V)$ for all $i = 1, 2, \dots, k$.*

Lemma 8 ([30, Lemma 3]). *If there is an epimorphism of G onto some group which doesn't have proper verbal subgroups of finite width and which is rich by verbal subgroups then G doesn't have proper verbal subgroups of finite width.*

Recall, that the braid group $B_n(\mathcal{G}_\emptyset^1)$ is Artin group of type B . From some results of the papers [29, 30] follows that the braid group $B_n(E^2) = B_n(\mathcal{G}_\emptyset^2)$ of Euclidean plane, the braid groups $B_n(\mathcal{G}_\emptyset^1)$, and corresponding pure braid groups $P_n(E^2)$ and $P_n(\mathcal{G}_\emptyset^1)$ for $n \geq 3$ doesn't have proper verbal subgroups of finite width.

We suppose that holds

Hypothesis. For every genetic code $\mathcal{G}$ there is a natural number $N = N(\mathcal{G})$ such that for all $n \geq N$ the braid group $B_n(\mathcal{G})$ doesn't have proper verbal subgroups of finite width.

For the pure braid group holds

Proposition 2. *Let $\mathcal{G} = \mathcal{G}_\emptyset^l, \mathcal{G}_0^l$ for $l \geq 2$ or $\mathcal{G} = \mathcal{G}_1^l$ for $l \geq 3$. Then for every $n \geq 1$ the pure braid group $P_n(\mathcal{G})$ doesn't have proper verbal subgroups of finite width.*

Proof. Let V be a finite proper set of words. We can see that for the pure braid group $P_n(\mathcal{G})$ there is an epimorphism on n -th direct power $H = G(\mathcal{G}) \times G(\mathcal{G}) \times \dots \times G(\mathcal{G})$. In view of the lemma 7 $\text{wid}(H, V) = \text{wid}(G(\mathcal{G}), V)$. The group $G(\mathcal{G})$ is either a free non-abelian or a group with one relation and ≥ 3 generators. In the both cases $G(\mathcal{G})$ is rich by verbal subgroups. The fact that $G(\mathcal{G})$ doesn't have proper verbal subgroups of finite width follows from [32, Theorem 2]. Hence, in view of the lemma 8 the group $P_n(\mathcal{G})$ also doesn't have proper verbal subgroups of finite width. The proposition is proven.

In connection with the defined construction we formulate the following questions.

Question 1. (I. A. Dynnikov). Let $\mathcal{G}$ be a genetic code with one defining relation, $K_{\mathcal{G}}$ be an one-cell complex that corresponding to the genetic code $K_{\mathcal{G}}$, $B_n(K_{\mathcal{G}})$ be the braid group on n strands, that corresponding to the complex $K_{\mathcal{G}}$ (it is defined similar to the braid group of manifold). In what cases the braid group $B_n(\mathcal{G})$ of the genetic code $\mathcal{G}$ is isomorphic to the braid group $B_n(K_{\mathcal{G}})$ of the complex $K_{\mathcal{G}}$?

Question 2. Let M be a two dimensions compact non-orientable manifold of genus g , $\mathcal{G} = \langle x_1, x_2, \dots, x_g \mid x_1^2 x_2^2 \dots x_g^2 = 1 \rangle$ be a genetic code that defines the fundamental group $\pi_1(M)$. How do the groups $B_n(M)$ and $B_n(\mathcal{G})$ connect for $n \geq 2$?

Question 3. Which groups $B_n(\mathcal{G})$ are decomposable by non-trivial manner in a free product with amalgamation?

REFERENCES

- [1] E. Brieskorn, K. Saito, Artin–Gruppen und Coxeter–Gruppen, *Invent. Math.*, 17 (1972), 245-271.
- [2] V. Ja. Lin, Artin braids and related groups and spaces, (Russian) *Algebra. Topology. Geometry*, Vol. 17, 159-227, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Informatsii, Moscow, 1979.
- [3] V. A. Vassiliev, Complements of discriminants of smooth maps: Topology and applications, *Translations of Mathematical Monographs*, vol. 98, Amer. Math. Soc., Providence, RI, 1992.
- [4] P. Cartier, Developpements recent sur les groupes de tresses. Applications a la topologie et a l’algebre. *Seminaire Bourbaki*, 1989-90, No 716, 1990, 17-68.
- [5] J. S. Birman, New points of view in knot theory, *Bull. Am. Math. Soc.*, New Ser. 28, No 2 (1993), 253-287.
- [6] L. H. Kauffman, Virtual knot theory, *Eur. J. Comb.*, 20, No 7 (1999), 663-690.
- [7] V. V. Vershinin, Braid groups and loop spaces. (Russian) *Uspekhi Mat. Nauk*, 54, No 2 (1999), 3-84; translation in *Russian Math. Surveys* 54, No 2 (1999), 273-350
- [8] R. Fenn, R. Rimányi, C. Rourke, The braid–permutation group, *Topology*, 36, No 1 (1997), 123-135.
- [9] V. G. Bardakov, The virtual and universal braids, *Fundamenta Math.*, 184, (2004), 1-18.
- [10] V. F. R. Jones, Hecke algebra representations of braid groups and link polynomials, *Ann. of Math.*, 126, No 2 (1987), 335-388.
- [11] S. Bigelow, Braid groups are linear, *J. Amer. Math. Soc.*, 14, No 2 (2001), 471-486.
- [12] D. Krammer, Braid groups are linear, *Annals of. Math.*, 155, No 1 (2002), 131-156.
- [13] P. Dehornoy, From large cardinals to braids via distributive algebra, *J. Knot Theory and its Ramifications*, 4, No 1 (1995), 33-79.
- [14] M. Atiyah, The Jones-Witten invariants of knots. *Seminaire Bourbaki*, 1989-90, No 715, 1990, 7-16.
- [15] M. Atiyah, The geometry and physics of knots. *Lezioni Lincee. [Lincei Lectures]* Cambridge University Press, Cambridge, 1990.
- [16] O. Zariski, The topological discriminant group of a riemann surface of genus p , *Amer. J. of Math.*, 59, No 2 (1937), 335-358.
- [17] G. P. Scott, Braids groups and the group of homeomorphisms of a surface, *Proc. Camb. Phil. Soc.*, 68, (1970), 605-617.
- [18] R. Gillette, J. Buskirk, The word problem and consequences for the braid groups and mapping class groups of the 2-sphere, *Trans. Amer. Math. Soc.*, 131, No 2 (1968), 277-296.
- [19] E. Fadell, J. Van Buskirk, The braid groups of E^2 and S^2 , *Duke. Math. J.*, 29, No 2 (1961), 243-258.
- [20] J. Van Buskirk, Braid groups of compact 2–manifolds with elements of finite order, *Trans. Amer. Math. Soc.*, 122, No 1 (1966), 81-97.
- [21] J. González-Meneses, New presentations of surface braid groups, *J. Knot Theory Ramifications*, 10, (2001), 431-451.
- [22] P. Bellingeri, On presentation of surface braids, *J. Algebra*, 274 (2004), 543-563.
- [23] P. Bellingeri, E. Godelle, Questions on surface braid groups, Preprint, Universite de Caen, CNRS UMR 6139, 2005-15, 13 p.
- [24] V. G. Bardakov, M. V. Neshchadim, Some property of braid groups of compact oriented 2-manifolds, IV International Algebraic Conference dedicated to the 60th anniversary of Professor Yu. I. Merzljakov, Novosibirsk, Institute of Mathematics, 2000, 9-13 (in Russin).
- [25] J. S. Birman, Braids, links and mapping class group, Princeton–Tokyo: Univ. press, 1974.
- [26] A. A. Markoff, Foundations of the algebraic theory of braids, *Trudy Mat. Inst. Steklova*, No. 16 (1945), 1-54.
- [27] M. I. Kargapolov, Yu. I. Merzljakov, *Fundamentals of the theory of groups*, Springer, New York, 1979.

- [28] H. S. M. Coxeter, W. O. J. Moser, Generators and relations for discrete groups. Fourth edition. *Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas]*, 14. Springer-Verlag, Berlin-New York, 1980.
- [29] V. G. Bardakov, On the theory of braid groups. (Russian) *Mat. Sb.* 183, No 6 (1992), 3-42; translation in *Russian Acad. Sci. Sb. Math.* 76 No 1 (1993), 123-153.
- [30] V. G. Bardakov, The width of verbal subgroups of some Artin groups. (Russian) *Group and metric properties of mappings (Russian)*, 8-18, Novosibirsk. Gos. Univ., Novosibirsk, 1995.
- [31] N. Burbaki, *Gruppy i algebrы Li. Gruppy Kokstera i sistemy Titsa. Gruppy, porozhdennye otrazheniyami. Sistemy kornei.* (Russian) [Lie groups and algebras. Coxeter groups and Tits systems. Groups generated by reflections. Root systems] Translated from the French by A. I. Kostrikin and A. N. Tjurin. Edited by A. I. Kostrikin. Izdat. "Mir", Moscow, 1972.
- [32] V. G. Bardakov, On the width of verbal subgroups of some free constructions. (Russian) *Algebra i Logika* 36 No 5 (1997), 494-517, 599; translation in *Algebra and Logic*, 36 No 5 (1997), 288-301

SOBOLEV INSTITUTE OF MATHEMATICS, NOVOSIBIRSK 630090, RUSSIA

E-mail address: bardakov@math.nsc.ru